



Vegetable Crop Update

A newsletter for commercial potato and vegetable growers prepared by the University of Wisconsin-Madison vegetable research and extension specialists

No. 9 – June 16, 2017

In This Issue

Late Blight and Early Blight Disease Forecast Updates
 National Late Blight Updates
 National Cucurbit Downy Mildew Updates
 Note from Dr. Yi Wang – New UW-Horticulture Vegetable Production Specialist

Calendar of Events

July 20, 2017 – UW-Hancock ARS Field Day, Hancock, WI
July 27, 2017 – UWEX Langlade County Airport Research Station Field Day, Antigo, WI
August 4, 2017 – UW-Lelah Starks Elite Foundation Seed Potato Farm Field Day, Rhinelander, WI (10AM to Noon Lunch to Follow)
January 21-23, 2018 – Wisconsin Fresh Fruit & Vegetable Conference, Wisconsin Dells, WI
February 6-8, 2018 – UWEX & WPVGA Grower Education Conference, Stevens Point, WI

Amanda J. Gevens, Associate Professor & Extension Vegetable Plant Pathologist, Interim Co-Director of Wisconsin Seed Potato Certification Program, UW-Madison, Dept. of Plant Pathology, 608-890-3072 (office), Email: gevens@wisc.edu. Webpage: www.plantpath.wisc.edu/wivegdis/

Current P-Day (Early Blight) and Severity Value (Late Blight) Accumulations (R.V. James, UW-Plant Pathology/R.V. James Designs): A P-Day value of ≥ 300 indicates the threshold for early blight risk and triggers preventative fungicide application. A DSV of ≥ 18 indicates the threshold for late blight risk and triggers preventative fungicide application. **Red** text in table below indicates threshold has been met/surpassed. “-” indicates that information is not available. Blitecast and P-Day values for actual potato field weather from Grand Marsh, Hancock, Plover, and Antigo are now posted at the UW Veg Path website at the tab “P-Days and Severity Values.”
http://www.plantpath.wisc.edu/wivegdis/contents_pages/weather_%20list_2017.html

Location	Planting Date	50% Emergence	P-Day Cumulative	Disease Severity Value	Date of DSV Generation	Increase in DSV from 6/10
<i>Antigo</i>	Early 5/3	5/25	104	17	6/16	14
	Mid 5/15	6/1	104	17	6/16	14
	Late 6/1	6/15	-	-	-	-
<i>Grand Marsh</i>	Early 4/10	5/15	191	20	6/16	9
	Mid 5/1	5/22	184	18	6/16	9
	Late 5/17	6/1	121	10	6/16	9
<i>Hancock</i>	Early 4/15	5/18	187	18	6/16	8
	Mid 5/5	5/30	124	8	6/16	8
	Late 5/20	6/5	86	8	6/16	8
<i>Plover</i>	Early 4/20	5/20	190	20	6/16	8
	Mid 5/8	5/25	165	9	6/16	8
	Late 5/25	6/8	69	8	6/16	8

Summary: Disease Severity Values (DSVs) and Late Blight Blitecast: All potatoes are at 50% emergence or greater. Several locations have reached threshold and should be considered for preventive fungicide application to manage the risk of late blight. Locations at/surpassing the threshold of DSV 18

are: early planted potatoes at Hancock, Grand Marsh and Plover; and mid-planted in Grand Marsh. We are nearly at threshold (DSV 17) in the Antigo area. Recall the maximum number of DSVs that one day can accumulate is 4. Once thresholds of 18 DSVs have been met, routine, protection of susceptible tomato and potato crops is recommended. Wisconsin commercial conventional fungicides for potato late blight control can be found at:

<http://www.plantpath.wisc.edu/wivegdis/pdf/2017/May%2022,%202017.pdf>

P-Days indicating early blight risk have not yet reached threshold for Wisconsin potatoes. Recall the threshold is 300 P-Days. In commercial fields planted in mid-April in southern/central Wisconsin, the first early blight lesions have been noted. We typically reach 300 P-Days on/around the first of July in the Hancock area for a general reference.

National Late Blight Updates: <http://usablight.org> is a useful resource for the detection and characterization of late blight on tomato and potato crops from the U.S. No new reports of late blight in the US have been reported at the site during recent weeks. However, reports of late blight were confirmed from parts of NC and VA last week. No clonal lineage/strain types have yet been described. Already this year, late blight has been confirmed on potato and tomato in southwestern Florida. In all reported cases, the pathogen genotype was US-23. This has been the predominant genotype in Wisconsin, and across the U.S., in recent years. US-23 can still generally be managed well with use of phenylamide fungicides.

National Cucurbit Downy Mildew Updates: <http://cdm.ipmpipe.org/> offers information on the detection and characterization of the cucurbit downy mildew pathogen from the U.S. (and often Canada). In this past week, confirmations of downy mildew have come from GA, NC, SC, and TX. Prior confirmations were from FL, GA, NC, and TX on a variety of cucurbits.

Yi Wang, Assistant Professor & Extension Potato and Vegetable Production Specialist, UW-Madison, Department of Horticulture, Cell Phone: 608-335-0933, Email: wang52@wisc.edu.

I am originally from China, but I lived in Madison for 7.5 years before moving to the west, so Madison is like my second hometown. Glad to be back, Badgers!

As I am starting my new potato and vegetable sustainable production extension program, I am open to conversations with our growers and industry to learn more. Please feel free to email, call, text, me, or schedule in-person visits with me so I can travel to your farm and have my feet on the ground. I will be at our summer field days in Antigo, Rhinelander, and Hancock this summer, so if you prefer, we can chat then. Look forward to working with all of you. Thank you.

The 2017 A3422 Commercial Vegetable Production in Wisconsin Guide is now available for 2017. As in past years, the guide can be downloaded for free (link below) or a hard copy can be purchased from UWEX Learning Store for \$10. <https://learningstore.uwex.edu/Assets/pdfs/A3422.pdf>